

# Studies on Tetraploidization in Watermelon (*Citrullus Lanatus* Thunb.)

T. Vardhini Kumari<sup>1</sup>, P. Ashok<sup>2</sup> and K. Sasikala<sup>3</sup>

<sup>1</sup>PG Scholar, HC&RI, Dr YSRHU V R Gudem 534 101, A.P

<sup>2</sup>Scientist (Hort.) AICRP on Tuber crops HRS, Dr YSRHU V R Gudem 534 101, A.P

<sup>3</sup>HC & RI, Dr YSRHU V R Gudem 534 101, A.P

E-mail: <sup>1</sup>vardhiniaphu@gmail.com, <sup>2</sup>hortashok@gmail.com, <sup>3</sup>sasiagr@gmail.com

**Abstract:** Seedless watermelon cultivars are in high demand by consumers not only because their fruits are seedless but also because they are sweeter than the fruits from diploid and seeded cultivars. Diploid watermelon varieties were treated with the chemical mutagen colchicine to develop tetraploid seed parents. Seedlings of different genetic back grounds (Arka Manik, Arka Muthu, Sugar Baby, IIHR-14) were raised and injected with different treatment combinations of colchicine (0.2%, 0.3%, 0.4%) with and without 0.5% PEG. Watermelon varieties responded differently for different treatment combinations of colchicine. The treated seedlings were evaluated for the tetraploidy induction on the basis of chloroplast count. The results revealed that the variety Arka Manik was sensitive to the colchicine treatments and among the four varieties, maximum number of tetraploids were identified in the variety Arka Muthu at 0.2 % colchicine, whereas 0.3 % colchicine+PEG was found to be the best for variety Arka Manik and 0.4 % colchicine+PEG, 0.4 % colchicine treatments were found to be best for the Sugar Baby and IIHR-14 respectively. The efficiency rate of tetraploid induction levels in the present experiment varied with the varieties selected even though they belong to same crop.

## 1. INTRODUCTION

Watermelon is a major cucurbit and an important crop in India. It is cultivated in an area of 76000 ha with an annual production of 1.70 MT in our country [7]. As a popular summer fruit and dessert, watermelon had been investigated continually to improve its eating quality. It is a fact that fruits contain plenty of seeds impairing the flesh quality. The presence of numerous seeds in the flesh disturbs the palate and charm of enjoying its sweet taste and granular texture. Seedless watermelon provides an opportunity to satisfy our sense of taste without fear of swallowing the seeds. Seedless watermelon cultivars are in high demand by consumers not only because their fruits are seedless but also because they are sweeter than the fruits from diploid and seeded cultivars [5]. The use of interploid hybridization between tetraploid and diploid plants has been the most effective method to obtain triploid seedless progeny.

Seedless watermelons have special virtues such as extra hard rind and firmer flesh which contribute to less damage in

shipment and longer shelf life for distant markets along with deep colour and highly stable soluble solids content [4]. Traditionally, tetraploid parents have been obtained by treating the newly emerged diploid seedlings with colchicine [10] and they were identified by counting the number of chloroplast per guard cell pair of fully expanded leaves [6]. However, lack of suitable tetraploid breeding parents has hampered progress towards the development of wide range of quality seedless cultivars [1]. Hence the present research was undertaken to standardize the colchicine concentration to induce of tetraploidy in four genetic backgrounds (Arka Manik, Arka Muthu, Sugar Baby and IIHR - 14).

## 2. MATERIAL AND METHODS

The present investigation was carried-out during *rabi* and *summer* seasons of 2012-2013 at the experimental plots of the Division of Vegetable Crops, Indian Institute of Horticultural Research (IIHR), Bangalore. Soils are red sandy loam with a pH ranging from 5.2 to 6.4. Four varieties of watermelon (Arka Manik, Arka Muthu, Sugar Baby and IIHR-14) possessing diverse agronomic and quality characters were selected for the study. The experiment was laid out in a factorial RBD with 32 treatment combinations replicated thrice with 100 plants per each replication. The seedlings were treated with different concentrations of colchicine at cotyledonary stage before the true leaf appeared. 20 µl of each treatment was applied to the growing tips of the seedlings for six consecutive days in the evening hours. Putative tetraploids were identified based on the chloroplast count in the guard cells and observing stomatal density per microscopic area [2]. Three concentrations of 0.2, 0.3 and 0.4 % of colchicine were tested for their efficacy in induction of tetraploidy. Poly Ethylene Glycol (PEG) is used as an adjuvant and expected to increase the permeability of colchicine into the cell wall. Hence the PEG at a concentration of 0.5 per cent was also used along with different concentrations of colchicine. The details of colchicine treatments and their preparations are mentioned in table 1.

**Table 1: Colchicine treatments and their preparations**

Concentration	Description
0.2% colchicines	0.2g of colchicine in 100 mL of distilled water
0.2% colchicine+0.5% PEG	0.2g of colchicine+0.5g of PEG in 100mL of distilled water
0.3% colchicines	0.3g of colchicine in 100 mL of distilled water
0.3% colchicine+0.5% PEG	0.3g of colchicine+0.5g of PEG in 100mL of distilled water
0.4% colchicines	0.4g of colchicine in 100 mL of distilled water
0.4% colchicine+0.5% PEG	0.4g of colchicine+0.5g of PEG in 100mL of distilled water
Distilled water	100mL of distilled water
0.5% PEG	0.5g of PEG in 100mL of distilled water

**3. RESULTS & DISCUSSION**

Four varieties of watermelon were subjected to different treatments of colchicine as described in the material and methods. The ploidy in watermelon was assessed by counting the number of chloroplasts per guard cell and observing stomatal density per microscopic area. The number of chloroplasts in each of guard cell pair ranged from 5 to 7 in diploids and 10-14 in tetraploids [3]. In certain cases the number of chloroplasts varied from 8 to 9 in each guard cell pair which might be due to mixoploidy. Based on this, among the four varieties, highest grand mean percentage of tetraploids (14.16 %) were identified in the variety Arka Muthu (Table 2) followed by Arka Manik (10.90), Sugar Baby (10.62) and IIHR-14 (9.76). Chloroplast count could be a useful tool to determine ploidy as variability in the number of chloroplasts within the guard cells is genetically related to the variability in the meristamatic cells [11].

**Table 2: Effect of colchicine concentrations on induction of tetraploidization among four varieties of watermelon**

PEG (P)	Varieties (V)											
	Arka Manik (V1)		Me an	Arka Muthu (V2)		Me an	Sugar Baby (V3)		Me an	IIHR-14 (V4)		Me an
	P0 : Co ntr ol	P1 : 0.5 % PEG		P0 : Co ntr ol	P1 : 0.5 % PEG		P0 : Co ntr ol	P1 : 0.5 % PEG		P0 : Co ntr ol	P1 : 0.5 % PEG	
C0 (Contr ol)	0.0 (0.28)	0.0 (0.28)	0.0 (0.28)	0.0 (0.28)	0.0 (0.28)	0.0 (0.28)	0.0 (0.28)	0.0 (0.28)	0.0 (0.28)	0.0 (0.28)	0.0 (0.28)	0.0 (0.28)

C1 (0.2% colchicine)	11.70 (19.9)	12.46 (20.6)	12.08 (20.25)	38.60 (38.3)	4.82 (12.3)	21.71 (25.3)	10.60 (18.9)	14.71 (22.5)	12.06 (20.7)	14.41 (22.2)	6.60 (14.7)	10.50 (18.45)
C2 (0.3% colchicine)	17.40 (24.6)	18.06 (25.1)	17.73 (24.85)	14.91 (22.7)	25.80 (30.5)	20.35 (26.6)	16.27 (23.7)	14.72 (22.5)	15.04 (23.1)	15.10 (22.9)	13.22 (21.2)	14.15 (22.05)
C3 (0.4% colchicine)	14.66 (22.5)	13.46 (21.5)	14.06 (22)	9.80 (18.2)	19.43 (26.1)	14.60 (22.15)	5.1 (13.4)	23.32 (28.8)	14.3 (21.1)	17.60 (24.7)	11.02 (19.3)	14.3 (22)
Mean	10.94 (16.82)	10.99 (16.87)	10.90 (16.87)	15.82 (19.87)	12.5 (17.29)	14.16 (17.07)	8.05 (14.52)	13.17 (18.6)	10.10 (17.2)	11.77 (17.52)	7.7 (13.87)	9.76 (13.6)

\*Figures in parenthesis indicates Arc sin transformed values

	S.E(m)±	CD (P =0.05)
Variety (V)	0.074	0.223
Concentration (C)	0.074	0.223
PEG (P)	0.052	0.158
VXC	0.148	0.446
CXP	0.105	0.315
VXP	0.105	0.315
VXCXP	0.21	0.630

The percentage of induction of tetraploids in different varieties was affected significantly at various concentrations of colchicine (with and without PEG). Maximum percentage of tetraploids (17.73%) in the variety Arka Manik and 15.45% in the variety Sugar Baby were obtained when treated with 0.3% of colchicine solution, whereas the induction percentage was 21.71% in the variety Arka Muthu, at 0.2% colchicine and for IIHR-14 it was 14.30% at 0.4% colchicine. Higher number of tetraploids was recorded at 0.2% of colchicine [3, 10].

Significantly higher percentage of tetraploids (38.60 %) was recorded in the variety Arka Muthu when treated at 0.2% colchicines without PEG, 18.06 % in the variety Arka Manik at 0.3% colchicine+PEG, 23.32 % in the variety Sugar Baby at 0.4% colchicine+PEG and 17.60 % in the variety IIHR-14 (Table 2) when treated with 0.4% colchicine, respectively. 30% of tetraploids were obtained at 0.3% colchicine when mixed with lanoline paste and applied at the cotyledonary stage [9]. The efficiency rate of tetraploid induction levels in the present experiment varied with the varieties selected.

These identified polyploids were subjected for stomatal count per microscopic area. It was observed that the number of stomata per microscopic area were significantly fewer in

tetraploids than in diploids in all the varieties studied. Among the tetraploids, the maximum number of stomata per microscopic area was observed in the variety IIHR-14 (44.41) followed by Arka Manik (43.06), Arka Muthu (42.30) and Sugar Baby (41.13). The stomatal density was found to be high in diploid plants (control) compared to mixoploid and tetraploids. The diploids of IIHR-14, Arka Manik, Arka Muthu and Sugar Baby were 62.10, 62.21, 63.30 and 64.32 respectively (Table 3). Besides this, it was generally observed that the stomata were bigger in tetraploids when compared to the diploids. Increasing ploidy often results in increased cell size and results in reduced frequency of stomata per unit leaf area in tetraploids when compared to the diploids [8].

**Table 3: Variation in stomatal density in polyploids of watermelon**

Ploidy	Arka Manik	Arka Muthu	Sugar Baby	IIHR-14
Tetraploid	43.06	42.30	41.13	44.41
Mixoploid	44.21	44.53	45.51	47.90
Diploid (Control)	62.21	63.30	64.32	62.10
CD (P=0.01)	3.90	4.60	3.80	2.70
S.E(m)±	1.30	1.50	1.20	0.90
CV (%)	11.02	13.30	10.7	10.99

#### 4. CONCLUSION

Based on the findings of the present investigation it could be concluded that variety Arka Manik was sensitive to the colchicine treatments. Among the four varieties, maximum number of tetraploids were identified in the variety Arka Muthu at 0.2 % colchicine, whereas 0.3 % colchicine+PEG was best for variety Arka Manik and 0.4 % colchicine+PEG, 0.4 % colchicine treatments were found to be best for the Sugar Baby and IIHR-14 respectively.

#### 5. ACKNOWLEDGEMENTS

This work was supported in part by a grant in the form of stipend from Dr Y. S. R. Horticultural University.

#### REFERENCES

- [1] Andrus, L., Ding, Z., Jiang, F., Jin, B.L.W., Ding, X., Sun, J., and Guiyuan, L., "Induction and Identification of Hexadecaploid of *Pinellia ternate*", *Journal Euphytica*, 1971, pp. 479-488.
- [2] Guimaraes, and Stotz, A., "Obtention of haploid plants induced by irradiated pollen in watermelon", *Cucurbit Genetics Cooperative*, 2004, pp.109-110.
- [3] Jaskani, M. J., Sung, W., Kwon, A., Dae, H., "Flow cytometry of DNA content of colchicine treated watermelon as a ploidy screening method", *Pakistan Journal of Botanicals*, 2005 pp. 685-696.
- [4] Kihara, H., and Nishiyama, A., "Triploid Watermelon", *Proceedings of American Society Horticulture Sciences*, 1951, pp. 217-230.
- [5] Marr, C. W., Gast, K. L. B., "Reactions by consuming in a farmers market to prices for seedless watermelon and rating of eating quality", *Hort Technology*, 1991, pp. 105-106.
- [6] McCuiston, F., and Elmstrom, G. W., "Identifying polyploids of various cucurbits by various guard cell chloroplast number", *Proc. Fla. State Hort. Society*, 2002, pp. 155-157.
- [7] National Horticulture Board. 2012 *Indian Horticulture Database* Gurgaon, New Delhi.
- [8] Nigel, M., R., Morpurgo, J., Dolezel, and Afza, R., "Induction and verification of autotetraploids in diploid banana (*Musa acuminata*) by *in vitro* techniques", *Euphytica* 2007, pp. 25-34.
- [9] Pitchaimuthu, M. O. P. D., Prasad, V. S. R. K., Swamy, K.R.M., "Development and Evaluation of Triploid Seedless Watermelon in India", *4<sup>th</sup> International Cucurbitaceous symposium*, 2004, pp. 82-83.
- [10] Suying, T., Xiuqiang, H., Liu, J., "A Study on the Increase of Induction Frequency of Tetraploid Watermelon", *Acta Agriculturae Boreali Sinica*, 1995, pp. 67-72.
- [11] Yudanova, B.B., and Zhang, X.P., "Hybrid seed production in watermelon", *Journal of New Seeds*, 2002, pp. 69-88.