International Conference on Contemporary Issues in Integrating Climate-The Emerging Areas of Agriculture, Horticulture, Biodiversity, Forestry; Engineering Technology, Fundamental/Applied Science and Business Management for Sustainable Development (AGROTECH-2017)

## Insect Infestation and Preventive Measures of Dried Fishery Products in Chalan Beel Area, Bangladesh

Syeda Nusrat Jahan<sup>1</sup>, Fouzia Nasrin<sup>2</sup>, Fawzia Adib Flowra<sup>3</sup>, Md. Bayezid Alam<sup>4</sup> and Saleha Jasmine<sup>5</sup>

<sup>1</sup>Department of Fisheries University of Rajshahi, Rajshahi-6205 Bangladesh <sup>2</sup>MS Student Department of Fisheries University of Rajshahi, Bangladesh <sup>3,4,5</sup>Department of Fisheries University of Rajshahi, Bangladesh E-mail: <sup>1</sup>nusratru@yahoo.com, <sup>3</sup>flowrabd@yahoo.com, <sup>4</sup>alambayezid@gmail.com, <sup>5</sup>psjasmine@yahoo.com

Abstract—The present study was conducted with regard to insect infestation and preventive measures to combat the blowfly infestation of the dried punti fish (Puntius sp.) in Chalan beel area, Bangladesh. In this study five treatments were done in sunny weather such as:  $T_1$  (without scaling, gutting and washing as control),  $T_2$  (with red chili powder),  $T_3$  (with turmeric powder),  $T_4$  (preserved with neem leaves) and  $T_5$  (with scaling, gutting, washing and net covering) from January 2015 to December 2015. Organoleptic test was carried out after six months storage period which indicated that  $T_5$  showed excellent result among all the treatments. In case of appearance, the highest mean value  $(8.06\pm0.94)$  was found in  $T_5$  and the lowest value (2.94±1.57) was found in  $T_1$ . With respect to colour, the highest mean value  $(7.50\pm0.71)$  and the lowest mean value  $(3.13\pm1.36)$ were found in  $T_5$  and  $T_1$  respectively. In terms of odour, the highest (8.06±0.78) and the lowest (2.88 $\pm$ 1.64) mean values were obtained in  $T_5$  and  $T_1$  respectively. In case of texture, the highest (8.06±0.78) and the lowest (2.88±1.64) mean values were observed in  $T_5$  and  $T_1$  respectively. The dried fishes treated with turmeric powder  $(T_3)$  exhibited good result having better quality whereas red chili  $(T_2)$  and neem leaves treated fishes  $(T_4)$  turned out to be acceptable in terms of quality.

Keywords: Dried fish, Insect infestation, preventive measures.

ISBN-978-93-85822-49-0

71