

Physico-Chemical and Microbial Characterization of Probiotic Plum Beverage

Jasia Nissar and Tehmeena Ahad

Sheri-kashmir University of Agricultural Sciences and Technology, Kashmir

Abstract—The plum juice with 13°Bx inoculated using 10% mixed culture of *L. bulgaricus* and *L. acidophilus* (1:1) and fermented for 7 hours was standardized for development of probiotic plum beverage. Various physico-chemical attributes of the beverage like TSS, titratable acidity, pH, total sugars, reducing sugars and non reducing sugars, ascorbic acid and polyphenol content were found to be 13 °Bx, 0.520 per cent, 3.512, 12.4 per cent, 11.61 per cent, 0.79 per cent, 8.57 mg/100mL and 283..21 mg/100mL, respectively. Further, viscosity of beverage was found to be decreased with increase in temperature. The microbiological analysis showed that prepared beverage contained optimum level of cultures i.e. 6.5×10^9 CFU/mL and was free from any traces of yeast, mold and coli-form bacteria. Hence, it can be concluded that the plum juice has the potential as a carrier of probiotic culture for the preparation of pomegranate probiotic beverage.

Keywords: Plum juice, Probiotics, Physico-chemical properties.

1. INTRODUCTION

Currently food industry is targeting for the development of more healthy foods due to consumers' awareness towards the relationship between food and health and their demands of decreasing the use of chemical preservatives. Nowadays, healthy foods mean "functional foods" which exert the beneficial effects on specific body functions, in addition to the traditional nutritional effects. Well-known examples of functional foods are those which contain bioactive compounds, like phytochemicals, oligosaccharides, dietary fiber and "friendly" bacteria i.e. probiotics⁷.

Probiotic foods are defined as those that contain microorganisms which influence beneficially the consumer's health by improving their intestinal microbial balance⁵. A commercial probiotic product is considered as functional only if it contains 10^7 CFU/ml at the time of consumption. Dairy is a very nutritious substrate for LAB but expanding the trend of vegan lifestyles; the issues of lactose intolerance and demand for low-fat and low-cholesterol foods have created a need for non-dairy probiotic products¹⁴.

Fresh fruits and vegetables are considered good matrices and can provide ideal substrates for probiotics and do not contain any dairy allergens⁹. Plum is a wonderful fruit having health-promoting properties with, anticancer, antioxidant and

antimutagenic effects. Plum juice is a good source of sugars, fiber, organic acids, phytonutrient, including antioxidants that play role in disease prevention.

A wide range of studies have been carried out to find the potential of fruit juices such as tomato, beet and cabbage juices as raw materials for the production of probiotic drinks. Results have shown that all the strains (*L. plantarum*, *L. acidophilus* and *L. delbrueckii* except *L. casei*) are capable of growth in the fruit juices mentioned^{22,23,24}.

Since, in addition to being delicious a, the plum juice may be an excellent medium for the supplementation of existing nutraceutical components with probiotic culture. It is, therefore, felt logical to seek its potentiality for developing probiotic apple beverage.

2. MATERIALS AND METHODS

Preparation of pomegranate juice

Freshly harvested plum fruits were procured from local market of Srinagar (kashmir). Plum juice was prepared by blending the fruit in the domestic mixer. Its total soluble solids was maintained to 13°Bx and stored at 4°C before use.

Probiotic strains

Lactobacillus isolates, *Lactobacillus bulgaricus* and *L. acidophilus* were selected and stored at -20 °C in sterile screw cap tubes. **Preparation of starter culture**

The starter culture was prepared with the help of method described by Mousavi *et al.*¹¹, with slight modifications. *L. acidophilus* and *L. bulgaricus* was cultivated separately in the MRS broth for 24-h at 37°C. To obtain the biomass, 10 mL of the separately cultivated MRS broths were mixed in equal proportion (1:1) and centrifuged at 4000 rpm for 10 min. The obtained biomass was washed with sterile saline solution twice to remove the residual MRS media. Thus, inoculum was prepared.

Preparation of probiotic pomegranate beverage

Above prepared starter culture (10mL) was then added to the pasteurized plum juice (100 mL) to obtain 10% inoculation. It

was allowed to ferment in incubator at 37°C for 7 h. After incubation, the beverage was kept at refrigeration temperature for future use. **Physico-chemical analysis of probiotic pomegranate beverage**

Following physico-chemical properties of fresh pomegranate juice were determined: **Total soluble solids (T.S.S.), Titratable acidity and pH** Total soluble solids were measured immediately after extraction using hand refractometer. Titratable acidity, expressed as per cent lactic acid, was determined by titration against 0.1N NaOH using phenolphthalein as an end point indicator. The pH value was obtained by using a digital pH meter.

Total Sugars, Reducing Sugar and Non- reducing sugars

Total carbohydrate/sugars was estimated by standard procedure using phenol sulphuric acid¹³. The amount of reducing sugar of fresh juice was calculated by Nelson – Somogyi method²⁰ and non-reducing sugar was obtained by subtracting reducing sugars from total sugars.

Ascorbic acid (vitamin C)

Ascorbic acid contents of samples were determined according to the titration method using 2, 6-dichlorophenol indophenols¹⁵.

Total phenolic content

The concentration of phenolic compounds was determined by the Folin-Ciocalteu colorimetric method¹⁸ where 5g of sample was homogenized in 25 mL of 50% (v/v) ethanol/water solution. The sample (100 µL) was mixed with 5 mL of the 0.2N Folin- Ciocalteu reagent and 4 mL of 7.5% sodium carbonate. The mixture was kept for 2 h at room temperature in the dark before the absorbance was measured at 765 nm spectrophotometrically. The total phenolic content was expressed as mg gallic acid equivalents (mg GAE/100 mL).

Consistency (Viscosity)

Viscosity was determined using the viscometer at constant speed of 100 rpm and varying temperature with a spindle no S-62²¹. Viscosity was expressed in terms of centipoises (cP). Parameters used for viscosity measurement of plum juice were as follows:

- Shear rate: 10
- Speed: 100 rpm
- Temperature: 20, 30, 40°C

Microbial analysis

The viable count of mixed culture was determined by the standard plate count method using Man-Rogosa-Sharpe agar (MRS agar) and the results were expressed as CFU ml⁻¹ juice. The yeast and mold count of beverage was determined using

potato dextrose agar medium. The coli-form and basically *E. coli* are the indicator microbes of water contamination by feces. The coli-form gives red pink color colonies on the MacConkey agar. Plates were incubated at 37°C for 48-72 hours³.

3. RESULTS AND DISCUSSION

Physico-chemical characteristics of probiotic pomegranate beverage

The chemical properties of the prepared product are presented in Table 1.

Table 1: Chemical characteristics of probiotic plum beverage

S. No.	Paramter(s)	Observations
1	Total Soluble Solids (□Brix)	13.0
2	Acidity (% of lactic acid)	0.520
3	pH	3.510
4	Total sugars (%)	12.4
5	Reducing sugars (%)	11.61
6	Non reducing sugars (%)	0.79
7	Ascorbic Acid (mg/100mL)	8.57
8	Total phenolic content (mg/100mL)	283.21

From the Table 1, it is revealed that the TSS of probiotic pomegranate beverage was found to 13 □Bx as it was kept fixed during the preparation of beverage. The titratable acidity is a measure of shelf life of the product and guard against the attack of micro-organisms. It also helps to ensure some chemical changes during preparation¹⁹ and storage⁸. The titratable acidity was 0.520 percent expressed in terms of lactic acid, produced during the metabolic activity of probiotic organisms. The pH is inversely proportional to the acidity of any medium. The pH value observed was 3.510..

The probiotic plum beverage contained 12.4, 11.61 and 0.79 percent of the total sugars, reducing sugars and non reducing sugars, respectively. Ascorbic acid, total phenol content which contribute to antioxidant property of plum juice were found to 8.57 mg/100mg and 283.21 mg/100mL, respectively.

Viscosity characteristics of plum beverage

The viscosity is an important characteristic to determine flow properties of a fluid.. In the present investigation, the effect of temperature on the viscosity of probiotic plum beverage was estimated. The data related to viscosity change is tabulated in Table 2.

Table 2 Effect of temperature on viscosity characteristics of probiotic plum beverage

S. No.	Temperature (°C)	Viscosity (cP)
1	20	8.2
2	30	7.6
3	40	3.4

- Each value is an average of three determinations

As shown in the above table, the viscosity changed significantly with the increase in temperature. At temperature 20°C, the viscosity of beverage was higher (8.2 cP) than that observed at 30°C temperature (7.6 cP). The viscosity observed at 40°C was found significantly lower (3.4cP) showing a declining trend as the temperature increased. The reason for viscosity reduction was that the heat causes the molecules to speed up as they bump and move around each other. Hence, more temperature means more movement of molecules and thus reducing their resistance to flow. According to Magerramov *et al.*¹⁰, the viscosity of tangerine and lemon juices monotonically decreases with the temperature.

Microbial analysis

The growth of undesirable organisms will spoil the product and may lead to food borne diseases affecting the healthy lives. Therefore, performing microbial analysis is mandatory in probiotic based products to assess their safety. The data related to microbiological analysis of probiotic beverage is tabulated in Table 3.

Table 3 Microbial analysis of probiotic plum beverage

S. No.	Parameters	Observations
1	Total plate count (CFU/mL)	6.5 x10 ⁹
2	Yeast and mold count (CFU/mL)	ND
3	Coli-form count (MPN/mL)	ND

-Each value is an average of three determinations ND: not detected

In the present work, the count of beneficial bacteria was detected as 6.5 x10⁹ CFU/mL of beverage. This count was in range for a product to be called as probiotic¹⁷.

On the other hand, the yeast and mold count and coli-form count was also determined. And they were not detected in the sample, which showed that the product was free of any pathogenic microbes and safe for consumption.

4. CONCLUSION

The pasteurized plum juice was inoculated with probiotic cultures (10%) of *L. bulgaricus* and *L. acidophilus* (1:1) and fermented for 7 h. Results showed that the chemical parameters were in sufficient amount for providing nutrition and bioactive components to consumer. viscosity profile of the product was in acceptable range. Microbiological analysis found that the beverage contained the desired level of probiotic cultures (10⁹CFU/mL) which is helpful for maintaining the health of gastro intestinal tract. Further, the prepared beverage didn't contain any traces of yeasts and molds and also coli-form bacteria, thus indicating that beverage is containing only health benefitting bacteria.

REFERENCES

- [1] Charalampopoulos, D., Pandiella, S.S. and Webb, C., Application of cereals and cereal components in functional foods: A review. *International Journal of Food Microbiology*, **79**: 131-141 (2002).
- [2] Chris, B., Paul, N. and Anthony, P.W., Food Microbiology and Laboratory Practices. Blackwell Publishing, State Avenue, USA (2006).
- [3] Dogahe, M.K., Khosravi-Darani, K., Tofighi, A., Dadgar, M. and Mortazavian, M.A., Effect of Process Variables on Survival of Bacteria in Probiotics Enriched Pomegranate Juice. *British Biotechnology Journal*, **5(1)**: 37-50 (2014).
- [4] Fuller, R., Probiotics in man and animals. *Journal of Applied Bacteriology*, **66**: 365–378 (1989).
- [5] Gil, M.I., Toma's-Barbera'n, F.A., Hess- Pierce, B., Holcroft, D.M. and Kader, A.A., Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *J. Agric. Food Chem.*, **48**: 4581-4589 (2000).
- [6] Jankovic, I., Sybesma, W., Phoithirath, P., Ananta, E. and Mercenier, A., Application of probiotics in food products-challenges and new approaches. *Curr. Opin. Biotechnol.*, **21**: 175-181 (2010). Langthasa, S., Processing and Preservation of Apple Pulp. Ph. D. Thesis, IARI, New Delhi (1999). Luckow, T. and Delahunty, C., Consumer acceptance of orange juice containing functional ingredients. *Food Research International*, **37**: 805-814 (2004).
- [7] Magerramov, M.A., Abdulagatov, A.I., Abdulagatov, I.M. and Azizov, N.D., Viscosity of tangerine and lemon juices as a function of temperature and concentration. *International Journal of Food Science and Technology*, **42**: 804–818 (2007).
- [8] Mousavi, Z.E., Mousavi, S.M., Djomeh Emam, Z. and Kiani, H., Fermentation of pomegranate juice by probiotic lactic acid bacteria. *World J Microbiol Biotechnol*, DOI 10.1007/s11274-010-0436-1 (2010).
- [9] Ranadheera, R.D.C.S., Baines, S.K. and Adams, M.C., Importance of food in probiotic efficacy. *Food Research International*, **43**: 1-7 (2010).
- [10] Ranganna, S., Handbook of Analysis and Quality Control for Fruits and Vegetable Products (2nd Edition). Tata McGraw-Hill Publishing Company Ltd., New Delhi (1991).
- [11] Saxena, V., Mishra, G., Saxena, A. and Vishwakarma, K.K., A comparative study on quantitative estimation of tannins in *Terminalia chebula*, *Terminalia belerica*, *Terminalia arjuna* and *Saraca indica* using spectrophotometer. *Asian J Pharm Clin Res*, **6 (3)**: 148-149 (2013).
- [12] Shah, N.P., Functional foods from probiotics and prebiotics. *Food Technology*, **55(11)**: 46-53 (2001).
- [13] Singleton, V.L. and Rossi, J.A.J., Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Amer. J. Enol. Viticult.*, **16**: 144-158 (1965).
- [14] Swientek, B., Toasts of the town. *Prep. Foods*, pp: 21-26 (1998).
- [15] Syed, H.M., Syed, I., Deshpande, H.W., Kulkarni, K.D. and Kulkarni, D.N., Chemical Analysis of Food Samples – Laboratory Manual. Needs Agencies, Parbhani (M.S.) (2007).

- [16] Weaver, M.C. and Daniel, R.J., *The Food Chemistry Laboratory - A Manual for Experimental Foods, Dietetics, and Food Scientists* (2nd Editions). CRC Press (2005).
- [17] Yoon, K.Y., Woodams, E.E. and Hang, Y.D., Probiotication of tomato juice by lactic acid bacteria. *J Microbiology*, **42(4)**: 315-318 (2004).
- [18] Yoon, K.Y., Woodams, E.E. and Hang, Y.D., Fermentation of beet juice by beneficial lactic acid bacteria. *Lebensm.-Wiss. u.-Technol.*, **38**: 73-75 (2005).
- [19] Yoon, K.Y., Woodams, E.E. and Hang, Y.D., Production of probiotic cabbage juice by lactic acid bacteria. *Bioresource Technology*, **97**: 1427-1430 (2006).